

NYXIA^{200mg} Tablets

Rifaximin

Nyxia tablets contains the active substance Rifaximin, an intestinal antibiotic used to treat: traveller's diarrhoea in adults when the diarrhoea is not accompanied by fever or blood in the stools, or 8 or more unformed (soft or liquid) stools in the last 24 hours. Rifaximin tablets are not recommended for use in children (aged less than 18 years). You must talk to a doctor if you do not feel better or if you feel worse after 3 days. What you need to know before you take Rifaximin tablets.

Do not take Rifaximin tablets.

If you are allergic (hypersensitive) to rifaximin, to similar types of antibiotics (such as rifampicin or rifabutin) or to any of the other ingredients of this medicine.

If you have a fever.

If you have blood in your stools.

If you passed 8 or more unformed stools in the last 24 hours.

If you have constipation, abdominal pain and vomiting caused by blockage of the bowel.

Warning and Precautions:

Talk to your doctor or pharmacist before taking Rifaximin tablets and take special care. If after 3 days of treatment, your symptoms continue or re-appear shortly afterwards do not take second course of Rifaximin tablet, see a doctor if your symptoms get worse during treatment stop taking Rifaximin tablets and consult a doctor. Other medicines and Rifaximin tablets. Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription. Please tell your doctor if you are taking ciclosporin (a medicine to suppress the body's immune system), warfarin (medicine to prevent blood clotting) or oral contraceptives (medicines to prevent pregnancy). If you are using activated charcoal (for example to treat wind or diarrhoea) please take Rifaximin tablets at least 2 hours after taking charcoal. Rifaximin tablets with food and drink Rifaximin tablets can be taken with or without food. Orally with a glass of water Taking this medicine may cause a reddish discolouration of your urine. Pregnancy, breast-feeding and fertility Nyxia tablets are not recommended during pregnancy or in fertile women not using contraception. Inform your doctor. If you are pregnant , think you may be pregnant or are thinking of becoming pregnant If you are breast-feeding or planning to start breast-feeding Ask your doctor or pharmacist for advice before taking any medicine. Driving and using machines Rifaximin tablets are unlikely to affect your reactions when driving or using machines. If you feel dizzy or drowsy you should not drive or operate machinery.

How to take Rifaximin tablets:

Always take Rifaximin tablets exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. Unless otherwise prescribed by the doctor, the usual dose is 1 tablet every 8 hours (600 mg/day). You should continue taking Rifaximin tablets for three day seven if your symptoms have improved. Unless otherwise prescribed, the duration of the treatment should not exceed three days. If your symptoms persist for more than three days please see a doctor. Do not break or crush the tablets.

If you take more Rifaximin tablets than you should contact your doctor or pharmacist.

If you take more than the recommended number of tablets, please contact a doctor.

If you forget to take Rifaximin tablets.

Take the missed dose as soon as you remember and take the next scheduled dose at its regular time.

If you stop taking Rifaximin tablets.

If you do not complete the three days of treatment recommended, your symptoms may worsen. If you have any further question on the use of this product, ask your doctor or pharmacist.

Possible Side Effects:

Like all medicines Rifaximin tablets can cause side effects, although not everybody gets them. If you have any of the following side effects stop taking Rifaximin tablets, and seek urgent medical advice immediately. These may be symptoms of a severe allergic reaction. Rash, hives or itchy skin. Shortness of breath, wheezing, Heart racing, Swelling of the mouth, face, ankles or any other part of the body. Extreme tiredness, dizziness or feeling faint. Common (may affect up to 1 in 10 people), Headache. Wind, abdominal bloating, abdominal pain, constipation, diarrhoea, urgency to empty, your bowels, nausea, involuntary and painful or ineffective straining, vomiting, fever. Uncommon (may affect up to 1 in 100 people) Thrush, cold sore, inflammation or infection of the nose and throat. Abnormal test results in white blood cells. Loss of appetite, loss of body fluid (dehydration). Abnormal dreams, depressed mood, sleeplessness, nervousness. Numbness, migraine, pins and needles, sinus headache, drowsiness. Double vision. earache, sensation of the room going round (vertigo) Increased blood pressure. Hot flushes. Cough, dry throat, blocked nose, sore throat, runny nose. upper abdominal pain, indigestion, intestinal movement disorder, dry lips, hard stools, blood in the stools, mucus in stools, taste disorders. Blood test results: increased liver enzyme values Sunburn. Back pain, muscle cramps, muscle weakness, muscle pain, neck pain. Abnormal urine test results: blood in urine, protein in urine, sugar in urine. Frequent urination, excessive urination. Frequent periods. Chills, cold sweat, increased sweating, flu-like illness, pain. The following side effects have been reported, however their frequency cannot be estimated from the available data: Bacterial infections Abnormal blood tests results (blood not clotting normally and abnormal liver tests). Allergic reactions to the drug. Skin roughness, skin redness, skin with little purple-coloured spots. Treatment with any antibiotic may cause Clostridium difficile associated diarrhoea (CDAD).

Reporting of side effects:

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

Storage:

Store between 15 to 30 °C.

Protect from heat, light and moisture.

Keep all medicines out of the reach of children.

Presentation:

Tablets are provided in a blister pack containing 1 x 10's tablets.

نگز یا 200 ملی گرام ٹیبلٹس
(ریفیکسیمین)

خاک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات: 15 سے 30 ڈگری سینٹی گریڈ تک رکھیں۔

گرمی، روشنی اور نمی سے بچائیں۔

تمام دوائیں بچوں کی پہنچ سے دور رکھیں۔

Manufactured by :
Pharmedic Laboratories (Pvt) Limited.
16 Km. Multan Road, Lahore - Pakistan